


# MELBOURNE NEUROSURGERY

## SURGICAL PROCEDURE INFORMATION

Prepared  
For

Dr. Damien Tange F.R.A.C.S.  
Neurosurgeon

## LUMBAR MICRO-LAMINECTOMY



16 Walpole St  
Kew Victoria 3101  
Australia  
[www.neurosurgery.com.au](http://www.neurosurgery.com.au)


## WHAT IS A MICRO-LAMINECTOMY ?

A Micro-Laminectomy is where part of the Lamina and part of the Facet Joint is removed to allow room for the Lumbar nerve. The nerve is usually compressed because of a degenerative process in the back.

## WHAT ARE THE REASONS FOR HAVING A MICRO-LAMINECTOMY ?

The commonest reason is because you are suffering from leg pain (called SCIATICA) you may also have numbness or weakness in your leg. If you are having surgery it usually means that the symptoms have not gone away with other therapies such as physiotherapy, rest, anti-inflammatories, nerve root blocks or epidurals. Most patients do get better with time but up to 15% may need surgery if the symptoms do not resolve. The cause of your symptoms is usually a degenerative process in the back where the facet joints enlarge, the disc bulges and the yellow ligament thickens. This all reduces the space for the nerves to the legs and causes your symptoms. Some times the surgery is done to try to improve any weakness or numbness that you may have in your leg.

A Micro-laminectomy is not normally performed for back pain alone as this generally does not improve with this type of surgery.



Picture of spine before the removal of the bone and ligament showing the compressed nerve root

## WILL ALL YOUR SYMPTOMS GET BETTER WITH SURGERY ?

This depends on the patient. Most will get substantially better and some will get complete resolution. This may depend on the amount of damage done to the nerve from the compression. Typically the leg pain is the first to improve. The muscle strength will then get better next but any thinning of the muscle may not improve. The strength does not always return back to normal. Numbness should improve but may not. It may take 9 months for this to be evident. Some patients will get increased low back pain after the surgery. Chronic pain may not improve.

## WILL YOU GET BACK TO NORMAL ?

Again this depends on the patient. Some will and some will not. The aim is to make a substantial improvement in your symptoms but some patients see no change at all. Not all your symptoms may be able to be treated with surgery and it may be that the micro-laminectomy is only planned to treat part of your problems.

## COULD YOU NEED FURTHER SURGERY ?

The spine is in a continual process of degeneration and the same surgery at the existing level may need to be repeated if you develop recurrent compression or scarring around the nerve. Surgery may need to be performed on the other side. Some patients may need more extensive surgery such as a laminectomy or fusion.

It is not unusual to need surgery at an adjacent level

## WILL THE SURGERY ALWAYS BE DONE AS DESCRIBED PRE-OPERATION ?

Some times while the surgery is being performed it may be felt that to get adequate control of the situation or to prevent further deterioration in the future that a different type of procedure may need to be performed. This may just be a further extension of the existing procedure or a much more involved procedure such as a spinal fusion. A smaller procedure may be done to reduce the risk of a later deterioration or because the operative findings are not as bad as suggested by your imaging.

## WHAT WILL HAPPEN IF YOU DO NOT HAVE THE PLANNED SURGERY ?


Generally the longer that you wait the more likely you are to get better so you may improve enough to not need the surgery.

The symptoms may not improve or may get worse.

If the compression worsens you may find that the symptoms of pain, numbness or weakness get worse.

If the compression gets really large you may develop compression of all the nerves to your bladder and bowel as well as the nerves to your legs. This produces paraplegia and loss of control of the bowel and bladder and is called a Cauda Equina syndrome and needs to be treated as an emergency.

Sometimes if the compression to the nerves is left too long before decompression it reduces the chances of a complete or any improvement following surgery.



### WHAT YOU NEED TO TELL THE DOCTOR BEFORE SURGERY ?

- If you have clotting problems.
- Any recent new Health problems or heart disease.
- If you are taking blood thinning agents.
  - e.g. Warfarin / aspirin/anti- inflammatory
- If you have improved from the time you decided to have surgery.
- Drug allergy
- Any thing that you think is important for the doctor to know in making his decision to offer you surgery.

### HOW LONG WILL YOU BE IN HOSPITAL ?

You may be admitted on the day of surgery or the day before. You will fast from midnight on the day of surgery. You will be discharged about 2- 3 days post-operatively. On discharge you will be able to perform most tasks of daily living (e.g. showering/dressing/etc.)

### HOW IS IT PERFORMED?


The surgery is performed in a hospital by a surgeon and he will most likely have a surgical assistant. In the operating theatre you are given a general anaesthetic by the anaesthetist (an indwelling catheter may be needed) and then positioned face down on a special frame. A small incision is marked out and the area prepared with antiseptic. You are covered in drapes so that only the incision can be seen.

A cut is made through the skin down to the spinous process. The muscle is dissected from the lamina / spinous process and facet joint and the retractor is used to hold this out of the way. The level is usually but not always checked with X-ray.


Using a special bone drill part of the lamina and the facet joint may be removed. This leaves the yellow ligament which is removed to expose the dura and the compressed nerve. The procedure may be needed on one or more nerves. It can be done bilaterally instead of a laminectomy leaving the spinous process intact.

Special attention is given to the nerves to make sure all pressure is removed. The openings under the facet joints that let the nerves out of the spine are checked and opened up if necessary.


Once this has been done all bleeding is stopped and a small piece of fat is placed behind the nerve to act as a cushion. The layers are then all sewn back to their normal positions. The skin may be closed with a nylon removable suture or with a dissolvable suture.



Picture of spine during the removal of the bone and before removal of prolapse using a highspeed drill



Picture of spine showing removal of the remaining yellow ligament under the facet joint



Picture of spine after closure showing the optional fat graft

### WHAT HAPPENS NEXT?

You will wake up in recovery and after about 1 hour you will be transferred to the ward. The nursing staff will be continually checking your pulse/blood pressure/and leg strengths looking for any changes to indicate a complication. During the first night on the ward you will be woken for these observations. You will have intravenous analgesia that you will control by pushing a button (this will be explained pre-op). Sometimes you will have difficulty passing water and you may require a catheter passed into your bladder. The catheter may be placed while you are asleep prior to starting the surgery.

The next day the drip in your arm will be removed after your first walk and then you will be given regular pain medication by mouth. Gradually over these next two days you will be able to get around as normal. When you are comfortable you will be able to go home. Some patients are not able to go directly home and will require either inpatient rehabilitation or to go home with a relative if you live alone.

It is important after the operation to walk as much as possible. Prolonged rest in bed can produce hip pain and clots in the legs.

Sometimes a couple of days after the operation the discomfort in your leg may return, this is due to swelling and usually settles with anti-inflammatory tablets.

If you have removable sutures then they are removed between 5 and 10 days.

### WHAT YOU SHOULD NOTIFY YOUR DOCTOR OF AFTER SURGERY ?

- Weakness in the legs
- Difficulty passing your urine
- Abdominal pain
- Increasing leg pain or numbness
- Fever
- Increasing back pain
- Swelling or infection in the wound.

### WHAT HAPPENS WHEN YOU GO HOME? (See the postoperative leaflet for more information)

You will be able to do most things.

You should avoid

- Heavy lifting
- Twisting
- Prolonged sitting

You will not be able to drive for 3 to 6 weeks

You should be able to return to some sort of work between 4 to 6 weeks.

It is important to walk as much as is comfortable.

If you smoke it is best to try to give this up.

If you are overweight then loss of weight is important

## **WHAT ARE THE RISKS?**

### **Discuss these and others with your surgeon**

Some of the possible complications are:

Infection (treated with antibiotics)

Damage to the nerve that is compressed

This may cause weakness, numbness and/or chronic pain

Damage to the dural sac containing the nerves and producing a fluid leak

Stops with bed rest but may need surgery to repair

Damage to structures in front of the disc

Bowel / ureter (tube from the kidney to the bladder) / Aorta (major blood vessel)

Post operative blood clot requiring drainage.

Wound breakdown

Paraplegia +/- loss of bowel and bladder function (very rare)

Impotence

Surgery on the wrong level or side

Blindness

Death

Disc prolapse at the level of surgery

Symptoms may be made worse

Further surgery may be required

Blood transfusion

Medical complications not related directly to the surgery

e.g. Clot in the legs (can travel to the lungs [uncommon])

Pneumonia

Heart attack

Urine infection

Kidney failure

Stroke

Drip or catheter infection

## **WHAT ARE THE COSTS OF THE SURGERY?**

### **Discuss this with your surgeon**

There are out of pocket costs for the surgery above the amount you will get back from your health fund. It is important to discuss this with your doctor and to contact your fund to understand not only the cost of the surgeon but also the costs of the hospital admission and other people involved. The assistant will send an account for 20% of the value of the surgeons' fee.

If you have no health insurance and wish to have the surgery privately then you should discuss with the office staff to organize an estimate.

YOU WILL BE ASKED TO SIGN A CONSENT FORM TO SAY THAT YOU UNDERSTAND ANY RISKS.

IF YOU ARE NOT SURE ASK BEFORE YOU SIGN.

If you feel that you need a second opinion we recommend that you ask your local doctor to arrange an independent opinion for you.

**MELBOURNE**  
**NEUROSURGERY**

[www.neurosurgery.com.au](http://www.neurosurgery.com.au)

PHONE 61-3-9 8532000  
FACSIMILE 61-3-9 8532500

MELBOURNE NEUROSURGERY  
PTY LTD  
ACN 082 289 316